

Return Service Requested



Ohio District 5 Area Agency on Aging, Inc. DECEMBER 2020 2131 Park Avenue West, Ontario, OH 44906 Caregiving in 419-524-4144 The Comfort of Home® **Caregiver Assistance News** CARING FOR YOU ... CARING FOR OTHERS"

Back Safety / Falls

Older adults are at risk of falling footwear and review because of the effects of medications, medications with his doctor. slowed reaction times, brittle bones, stiffness, and lack of flexibility and Fear of Falling impaired vision. Due to changes in the A person with AD, brain that are caused by Alzheimer's even in the early disease (AD), people with AD are at stages, will have especially high risk of falling. Slower subtle changes in reaction time, difficulty recognizing walking ability changes in the height or depth of that will become a step, for example, can lead to more severe as time tripping and falling. Changes in goes on. This can create difficulty balance and coordination combined with balance. If the person has with poor memory can make it other illnesses, the problems may be difficult for a person with AD to more severe. both get from one place to another and avoid hazardous objects at the It is natural that he will fear same time. He may miss a step while falling. To help the person in your looking for a door or trying to listen care feel more confident, adaptive to someone's conversation. You can devices such as walkers or canes can reduce the risk of falling by modifying be useful, but you will have to remind the environment. Try to help the him to use these devices. Bring the person in your care remain as active cane or walker to the person when he and flexible as possible. It is also has forgotten it. Exercises that you important to provide appropriate can do with the person in your care

Fall Facts

- ✓ Every second of every day, an older adult (age 65+) suffers a fall in the U.S.—making falls the leading cause of injury and injury death in this age group.
- Thirty million older adults fall each year—resulting in about 30,000 deaths. ~
- Each year, 3 million older adults are treated for a fall injury and at least ~ 300,000 seniors are hospitalized for hip fractures.
- Women fall more often than men and account for three-quarters of all 1 hip fractures.

```
Source: CDC.gov
```

Caregiving in The Comfort of Home®

Our Purpose

To provide caregivers with critical information enabling them to do their job with confidence, pride, and competence.

Ordering Info

From the publishers of

Caregiving in The Comfort of Home® **Caregiver Series**

available from ...

CareTrust Publications LLC PO Box 10283, Portland, OR 97296 800-565-1533 or www.comfortofhome.com

Comments and suggestions welcome.

©2020 CareTrust Publications LLC. All rights reserved. Reproduction of any component of this publication is forbidden without a license from the publisher

Some content in this publication is excerpted from The Comfort of Home: Caregivers Series. It is for informational use and not health advice. It is not meant to replace medical care but to supplement it. The publisher assumes no liability with respect to the accuracy, completeness or application of information presented or the reader's misunderstanding of the text

SAFETY TIPS-If the Person in Your Care Falls

When you suspect a broken bone, follow these steps:

- If the person cannot move or use the injured limb, keep it from moving. Do not straighten a deformed arm or leg.
- Support the injured part above and below the site of the injury by using folded towels, blankets, pillows, or magazines if the person cooperates.
- If the person is face down, and breathing is adequate, leave the person in the same position.
- Keep the person warm with a blanket and make the person as comfortable as possible.
- Call 911.

Source: Caregiving in the Comfort of Home: A Complete Guide for Caregivers



Article continues

on page 2

may improve his balance. Remember, before starting any type of exercise routine, get advice from your physician. Start slowly with only moderate effort. Give him or her time to build strength and stamina. Any amount of exercise helps reduce the risk of falls.

Back Safety - Proper Lifting and Moving

Back injuries are among the most common injuries in caregiving. But sometimes as a caregiver, you will have to help the person in your care move to a chair or be asked to pick up a heavy object. Though this seems like a simple maneuver, it is very easy to be injured doing this and left unable to perform your caregiving duties. A little attention to the safest way to move your body—and the person in your care—will keep you from unnecessary



injury. Most muscle strain injuries to both the caregiver and care receiver are preventable if you follow some simple rules.

Body Mechanics

The proper use of our bodies is called *body mechanics*. It involves standing and moving in ways that prevent injury, avoid fatigue, and make the best use of strength. When you learn how to move, control and balance your own body, it's easier to control and help or move another person.

General Rules

The following hints will help you use proper body mechanics and be safe. However, caregivers should seek training from a physical therapist to reduce the risk of injury.

- ➡ Only lift as much as you can comfortably handle.
- → Always let the person you are helping know what you are going to do.
- ➡ To create a base of support, stand with your feet 8"-12" apart with one foot a half step ahead of the other.
- ➡ Bend your knees slightly.
- ► Keep your spine in a neutral (normal arched curve, not stiff) position while lifting.
- ➡ USE YOUR LEGS instead of your back to do most of the work—leg muscles are stronger than back muscles.
- → If you can adjust the bed, keep the top at about waist level. If it is low or you are tall, put one foot on a footstool to relieve pressure on your lower back.

✓ Good nutrition. Physical fitness. ✓ Good body mechanics. ✓ A back strengthening exercise program than normal.

